

New Jersey Division of HIV/AIDS Services  
Persons Living with HIV/AIDS as of June 30, 2004  
Piscataway Residents

1. Current HIV/AIDS Category	Adult/Adolescent *		Pediatric *		Total	
	Cases	%	Cases	%	Cases	%
-----	-----	---	-----	---	-----	---
AIDS-Any OI	#	#	#	#	27	23
AIDS-CD4 Only	38	34	0	0	38	33
HIV Only	#	#	#	#	50	43
-----	-----	---	-----	---	-----	---
Total	112	100	#	#	115	100

Current			3. Race/Eth.	Adult/Adl*		Pediatric*		Total	
Age *	Cases	%		Cases	%	Cases	%	Cases	%
-----	-----	---	-----	-----	---	-----	---	-----	---
<5	0	0	White, Not Hispanic	35	31	0	0	35	30
5-12	#	#	Black, Not Hispanic	#	#	#	#	58	50
13-19	#	#	Hispanic	13	12	0	0	13	11
20-29	5	4	Other/Unknown	#	#	#	#	9	8
30-39	23	20	-----	-----	---	-----	---	-----	---
40-49	58	50	Total	112	100	#	#	115	100
50+	25	22							
Unk.	0	0							
-----	-----	---							
Total	115	100							

4. Patient Group	Adult/Adolescent Transmission Modes **					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Men sex with men (MSM)	25	34	0	0	25	23
Injection drug user (IDU)	20	27	11	30	31	28
MSM/IDU	#	#	0	0	#	#
Heterosexual contact	#	#	#	#	#	#
Risk not reported/other***	21	29	12	32	33	30
-----	-----	---	-----	---	-----	---
Total	73	100	37	100	110	100

	Pediatric Transmission Modes					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Parent at risk/has AIDS/HIV	#	#	#	#	#	#
Risk not reported/other***	0	0	#	#	#	#
-----	-----	---	-----	---	-----	---
Total	#	#	#	#	5	100

NOTE: Includes only those persons known to be infected with HIV.

# Indicates that number is not shown due to small cell size,  
in accordance with NJDHSS security and confidentiality policies.

\* Classification based on current age at the end of June 2004.

\*\* # patients are currently of adult/adolescent age (13+ years old), but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1,2, and 3, and as pediatric reports in table 4.

\*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services  
AIDS Cases Reported as of June 30, 2004  
Piscataway Residents

1. Disease Category	Adult/Adolescent*		Deaths (%)	Pediatric*		Deaths (%)	Total	
	Cases	(%)		Cases	(%)		Cases	(%)
PCP	60	( 34)	52 ( 87)	0	( 0)	0 ( .)	60	( 33)
Other OI w/o PCP	#	( #)	# ( #)	#	( #)	# ( #)	52	( 29)
KS Alone	#	( #)	# ( #)	#	( #)	# ( #)	7	( 4)
No Disease Listed	61	( 35)	23 ( 40)	0	( 0)	0 ( .)	61	( 34)
Total	175	(100)	113 ( 66)	5	(100)	# ( #)	180	(100)

2. Age *	Cases	%	3. Race/Eth.	Adult/Adl* Cases	%	Pediatric* Cases	%	Total Cases	%
Under 5	#	#	White, Not Hispanic	#	#	#	#	76	42
5-12	#	#	Black, Not Hispanic	#	#	#	#	90	50
13-19	#	#	Hispanic	8	5	0	0	8	4
20-29	21	12	Other/Unknown	6	3	0	0	6	3
30-39	81	45	-----	-----	---	-----	---	-----	---
40-49	49	27	Total	175	100	5	100	180	100
Over 49	23	13							
Unknown	0	0							
Total	180	100							

4. Patient Group	Adult/Adolescent Transmission Modes **				Total	%
	Males	%	Females	%		
Men sex with men (MSM)	48	35	0	0	48	27
Injection drug user (IDU)	55	40	14	38	69	39
MSM/IDU	#	#	0	0	#	#
Heterosexual contact	#	#	#	#	#	#
Risk not reported/other***	25	18	6	16	31	18
Total	138	100	37	100	175	100

	Pediatric Transmission Modes				Total	%
	Males	%	Females	%		
Parent at risk/has AIDS/HIV	#	#	#	#	5	100
Risk not reported/other***	0	0	0	0	0	0
Total	#	#	#	#	5	100

# indicates that number is not shown due to small cell size,  
in accordance with NJDHSS security and confidentiality policies.

\* Classification at time of AIDS diagnosis.

\*\* 0 patients were diagnosed with AIDS as adults, but have evidence either of having been HIV-infected or of having acquired AIDS as children. They are counted as adult/adolescent reports in tables 1,2, and 3; and as pediatric reports in table 4.

\*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services  
AIDS Cases Reported as of June 30, 2004  
Piscataway Residents

5. AIDS Cases by Year of Diagnosis.

Year of Diagnosis -----	Number of Cases -----	Number of Deaths -----	Case-Fatality Rate -----
Before 1985	#	#	#
1986	5	5	100%
1987	#	#	#
1988	15	14	93%
1989	8	5	63%
1990	10	10	100%
1991	18	14	78%
1992	20	19	95%
1993	19	14	74%
1994	16	10	63%
1995	20	8	40%
1996	14	#	#
1997	5	0	0%
1998	#	#	#
1999	#	#	#
2000	#	#	#
2001	7	#	#
2002	#	#	#
2003	#	#	#
-----	-----	-----	-----
Total	180	115	64%

New Jersey Division of HIV/AIDS Services  
HIV INFECTION-not-AIDS Reports with Identifiers Reported as of June 30, 2004  
Piscataway Residents

1. Age *	Cases	%	2. Race/Ethnicity	Adult/Adl* Cases	%	Pediatric* Cases	%	Total Cases	%
-----	-----	---	-----	-----	---	-----	---	-----	---
Under 5	#	#	White, Not Hispanic	15	27	0	0	15	27
5-12	#	#	Black, Not Hispanic	29	53	0	0	29	52
13-19	0	0	Hispanic	7	13	0	0	7	13
20-29	11	19	Other/Unknown	#	#	#	#	6	11
30-39	26	46	-----	-----	---	-----	---	-----	---
40-49	12	21	Total	55	100	#	#	57	100
Over 49	6	11							
Unknown	0	0							
-----	-----	---							
Total	57	100							

3. Patient Group	Adult/Adolescent Transmission Modes **					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Men sex with men (MSM)	14	41	0	0	14	25
Injection drug user (IDU)	7	21	6	29	13	24
MSM/IDU	#	#	0	0	#	#
Heterosexual contact	#	#	#	#	#	#
Risk not reported/other***	10	29	10	48	20	36
-----	-----	---	-----	---	-----	---
Total	34	100	21	100	55	100

	Pediatric Transmission Modes					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Parent at risk/has AIDS/HIV	0	.	0	0	0	0
Risk not reported/other***	0	.	#	#	#	#
-----	-----	---	-----	---	-----	---
Total	0	.	#	100	#	100

\* Classification at time of first HIV+ test or at time of report if not yet confirmed HIV+.

# indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.

\*\* 0 patients were diagnosed with HIV as adults, but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1 and 2; and as pediatric reports in table 3.

\*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

NOTE: These tables include those reports of HIV Positive infection that have not yet been updated to AIDS status. Not included in these tables are cases initially reported as HIV-not-AIDS that have subsequently been updated to AIDS. Also not included in these tables are HIV Perinatal Exposure Reports where the serostatus of the child is not confirmed HIV Positive; i.e., the HIV serostatus of the child is undetermined or has been found to be negative (seroreverted).

New Jersey Division of HIV/AIDS Services  
HIV/AIDS Cases Reported as of June 30, 2004  
Piscataway Residents

1. Current HIV/AIDS Category	Adult/Adolescent *		Pediatric *		Total	
	Cases	%	Cases	%	Cases	%
-----	-----	---	-----	---	-----	---
AIDS-Any OI	#	#	#	#	119	50
AIDS-CD4 Only	61	27	0	0	61	26
HIV Only	#	#	#	#	57	24
-----	-----	---	-----	---	-----	---
Total	230	100	7	100	237	100

2. Age *	Cases	%	3. Race/Eth.	Adult/Adl*		Pediatric*		Total	
-----	-----	---	-----	Cases	%	Cases	%	Cases	%
Under 5	#	#	White, Not Hispanic	#	#	#	#	91	38
5-12	#	#	Black, Not Hispanic	#	#	#	#	119	50
13-19	#	#	Hispanic	15	7	0	0	15	6
20-29	40	17	Other/Unknown	#	#	#	#	12	5
30-39	105	44	-----	-----	---	-----	---	-----	---
40-49	53	22	Total	230	100	7	100	237	100
Over 49	29	12							
Unknown	0	0							
-----	-----	---							
Total	237	100							

4. Patient Group	Adult/Adolescent Transmission Modes **					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Men sex with men (MSM)	62	36	0	0	62	27
Injection drug user (IDU)	62	36	20	34	82	36
MSM/IDU	5	3	0	0	5	2
Heterosexual contact	8	5	22	38	30	13
Risk not reported/other***	35	20	16	28	51	22
-----	-----	---	-----	---	-----	---
Total	172	100	58	100	230	100

	Pediatric Transmission Modes				Total	%
	Males	%	Females	%		
-----	-----	---	-----	---	-----	---
Parent at risk/has AIDS/HIV	#	#	#	#	5	71
Risk not reported/other***	0	0	#	#	#	#
-----	-----	---	-----	---	-----	---
Total	#	#	#	#	7	100

NOTE: Includes only those persons known to be infected with HIV.

# Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.

\* Classification at time of first HIV+/AIDS diagnosis.

\*\* 0 patients were diagnosed with HIV/AIDS as adults, but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1,2 and 3; and as pediatric reports in table 4.

\*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services  
HIV/AIDS Cases Reported as of June 30, 2004  
Piscataway Residents

5. HIV/AIDS Cases by Year of First HIV/AIDS Diagnosis.

Year of First HIV/AIDS Diagnosis	Number of Cases	Number Known Dead	Case-Fatality Rate
-----	-----	-----	-----
Before 1985	#	#	#
1985	#	#	#
1986	8	6	75%
1987	7	#	#
1988	18	17	94%
1989	17	10	59%
1990	16	13	81%
1991	26	17	65%
1992	22	12	55%
1993	17	12	71%
1994	18	8	44%
1995	17	9	53%
1996	15	#	#
1997	10	0	0%
1998	5	#	#
1999	6	#	#
2000	8	#	#
2001	7	#	#
2002	7	#	#
2003	5	0	0%
2004	#	#	#
-----	-----	-----	-----
Total	237	122	51%